

The entanglement dynamics of bipartite quantum system: Towards entanglement sudden death

Wei Cui^{1,2}, Zairong Xi^{1*} and Yu Pan^{1,2}

¹Key Laboratory of Systems and Control, Institute of Systems Science, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, People's Republic of China

²Graduate University of Chinese Academy of Sciences, Beijing 100039, People's Republic of China

E-mail: zrxixi@iss.ac.cn

Abstract. We investigate the entanglement dynamics of bipartite quantum system between two qubits with the dissipative environment. We begin with the standard Markovian master equation in the Lindblad form and the initial state which is prepared in the extended Werner-like state: $\rho_{AB}^{\Phi}(0)$. We examine the conditions for entanglement sudden death (ESD) and calculate the corresponding ESD time by the Wootters' concurrence. We observe that ESD is determined by the parameters like the mean occupation number of the environment N , amount of initial entanglement α , and the purity r . For $N = 0$, we get the analytical expression of both ESD condition and ESD time. For $N > 0$ we give a theoretical analysis that ESD always occurs, and simulate the concurrence as a function of $\gamma_0 t$ and one of the parameters N, α , and r .

PACS numbers: 03.65.Ud, 03.65.Yz, 03.67.Mn, 05.40.Ca

1. Introduction

Entanglement is responsible for the most counterintuitive aspects of quantum mechanics [1], and motivated many philosophical discussions in the early days of quantum physics [2]. Recently it has been regarded as a resource for quantum information processing [3]. In fact, entanglement is one of the key ingredient for quantum teleportation [4, 5, 6], quantum cryptography [7] and is believed to be the origin of the power of quantum computers, etc. However, a quantum system used in quantum information processing inevitably interacts with the surrounding environments (or the thermal reservoirs), which takes the pure state of the quantum system into a mixed state [8]. Thus, it is an important subject analyzing the entanglement decay induced by the unavoidable interaction of the interested systems with the environment [9]. In one-party quantum system, this process is called decoherence, and various methods have been proposed to reduce this unexpected effect [10, 11, 12, 13]. In multiparty systems with non-local quantum correlations much interest has been arisen in the dynamics of entanglement. For example, entanglement sudden death (ESD), which means that it disappears at finite time, was discovered by Yu and Eberly [14, 15]. It differs remarkably from the single qubit coherent evolution. The interesting phenomenon has been experimentally observed for entanglement photon pairs [16] and atomic ensembles [17].

ESD puts a limitation on the time when entanglement must be exploited. The evolution of the entanglement and ESD have been analyzed and various interesting results obtained [18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28]. Ref. [23] investigated the time evolution of entanglement of various entangled states of a two-qubit system. For four different initial states they analyzed the entanglement sudden death conditions and time. From the simulation they got the conclusion that the ESD always exists except for the vacuum reservoir. However, it is still a question when the ESD occurs and what ESD time is in the general cases and universal initial states. The aim of this paper is to discuss the above problems for the standard Markovian master equation in the Lindblad form and the initial states, the extended Werner-like states: $\rho_{AB}^\Phi(0)$.

The paper is organized as follows. We first introduce the Wootters' concurrence [29] and the extended Werner-like states: $\rho_{AB}^\Phi(0)$. In Section III we give a standard Markovian master equation [30, 31]. The master equation is equivalent to a first order coupled differential equations when the initial state is the extended Werner like state $\rho_{AB}^\Phi(0)$. In sections IV, we analyze the ESD conditions and ESD time for the initial state $\rho_{AB}^\Phi(0)$. Conclusions and prospective views are given in Section V.

2. Concurrence and initial states

A useful measure of entanglement is the Wootters' concurrence [29]. For a bipartite system described by the density matrix ρ , the concurrence $\mathcal{C}(\rho)$ is

$$\mathcal{C}(\rho) = \max(0, \sqrt{\lambda_1} - \sqrt{\lambda_2} - \sqrt{\lambda_3} - \sqrt{\lambda_4}), \quad (1)$$

where $\lambda_1, \lambda_2, \lambda_3$, and λ_4 are the eigenvalues (with λ_1 the largest one) of the “spin-flipped” density operator ζ , and

$$\zeta = \rho(\sigma_y^A \otimes \sigma_y^B) \rho^*(\sigma_y^A \otimes \sigma_y^B), \quad (2)$$

where ρ^* denotes the complex conjugate of ρ and σ_y is the usual Pauli matrix. $\mathcal{C}$ ranges in magnitude from 0 for a disentanglement state to 1 for a maximally entanglement state.

The general structure of an “X” density matrix [14, 15] is as follows

$$\hat{\rho} = \begin{pmatrix} x & 0 & 0 & v \\ 0 & y & u & 0 \\ 0 & u^* & z & 0 \\ v^* & 0 & 0 & w \end{pmatrix}, \quad (3)$$

with x, y, z, w real positive and u, v complex quantities. Such states are general enough to include states such as the Werner states, the Bell states, *et al.*. A remarkable aspect of the “X” states is that the initial “X” structure is maintained during the Lindblad master equation evolution. This particular form of the density matrix allows us to analytically express the concurrence at time t as [14]

$$\mathcal{C}_\rho^X(t) = 2 \max\{0, |u| - \sqrt{xw}, |v| - \sqrt{yz}\}. \quad (4)$$

In the present work, we will analyze in detail the two-qubit entanglement dynamics in Markovian environment starting from the initial “X” states defined in Eq. (3). We will examine the exactly ESD time and entanglement evolution for two kind of special states, the extended Werner-like states,

$$\rho_{AB}^\Phi(0) = r|\Phi\rangle_{ABAB}\langle\Phi| + \frac{1-r}{4}\mathbf{I}_{AB}, \quad (5)$$

where r the purity of the initial states, $\mathbf{I}_{AB}$ the 4×4 identity matrix and

$$|\Phi\rangle_{AB} = (\cos(\alpha)|10\rangle + \sin(\alpha)|01\rangle)_{AB}, \quad (6)$$

with α measuring the amount of initial entanglement. The state $\rho_{AB}^\Phi(0)$ has the following form

$$\rho_{AB}^\Phi(0) = \begin{pmatrix} \frac{1-r}{4} & 0 & 0 & 0 \\ 0 & r \cos^2(\alpha) + \frac{1-r}{4} & r \sin(\alpha) \cos(\alpha) & 0 \\ 0 & r \sin(\alpha) \cos(\alpha) & r \sin^2(\alpha) + \frac{1-r}{4} & 0 \\ 0 & 0 & 0 & \frac{1-r}{4} \end{pmatrix}. \quad (7)$$

Obviously, the state in Eq. (5) is reduced to the standard Werner state when $\alpha = \pi/4$, and the Werner-like state become totally mixed state for $r = 0$, while the well-known Bell state for $r = 1$. Note that, the extended Werner-like state $\rho_{AB}^\Phi(0)$ contain both separate state and entangled state. According to Peres’ criterion [32], when $\frac{1}{1+2|\sin 2\alpha|} \leq r \leq 1$ the extended Werner-like state would be entangled, otherwise it would be separated. Fig.1 shows the entangled and separated areas.




Figure 1. (Color online) Plot the initial entangled area (colored) of the extended Werner-like states for $\alpha \in [0, 2\pi]$. The blue line is $r = \frac{1}{1+2|\sin(2\alpha)|}$.

3. The master equation

The standard two-qubit Markovian master equation is the following Lindblad form [8, 30, 31],

$$\frac{d\rho}{dt} = \frac{\gamma_0(N+1)}{2} \sum_{j=1}^2 \{2\sigma_j^- \rho \sigma_j^+ - \sigma_j^+ \sigma_j^- \rho - \rho \sigma_j^+ \sigma_j^-\} + \frac{\gamma_0 N}{2} \sum_{j=1}^2 \{2\sigma_j^+ \rho \sigma_j^- - \sigma_j^- \sigma_j^+ \rho - \rho \sigma_j^- \sigma_j^+\}, \quad (8)$$

where $N = 1/(e^{\frac{\hbar\omega_0}{K_B T}} - 1)$, the mean occupation number of the environment oscillators, γ_0 is the spontaneous emission rate. As introduced before, we assume the initial state to be a “X” state. When substituting (3) into the master equation (8) we obtain the following first-order coupled differential equations,

$$\begin{pmatrix} \dot{x}(t) \\ \dot{y}(t) \\ \dot{z}(t) \\ \dot{w}(t) \end{pmatrix} = \begin{pmatrix} -2\gamma_0(N+1) & \gamma_0 N & \gamma_0 N & 0 \\ \gamma_0(N+1) & -\gamma_0(2N+1) & 0 & \gamma_0 N \\ \gamma_0(N+1) & 0 & -\gamma_0(2N+1) & \gamma_0 N \\ 0 & \gamma_0(N+1) & \gamma_0(N+1) & -2\gamma_0 N \end{pmatrix} \begin{pmatrix} x(t) \\ y(t) \\ z(t) \\ w(t) \end{pmatrix}, \quad (9)$$

and

$$\begin{aligned} \dot{u}(t) &= -(1+2N)\gamma_0 u(t), \\ \dot{v}(t) &= -(1+2N)\gamma_0 v(t). \end{aligned} \quad (10)$$

The solution of the previous master equation can be found by solving the system of differential equations. The reduced density matrix elements $x(t), y(t), z(t), w(t), u(t)$ and $v(t)$ are given

(i) $N > 0$

$$\begin{aligned}
 x(t) &= c_1 \frac{N}{N+1} - c_2 \Gamma^2(t) - c_4 \gamma_0 N \Gamma(t), \\
 y(t) &= c_1 + c_2 \Gamma^2(t) + c_3 \Gamma(t), \\
 z(t) &= c_1 + c_2 \Gamma^2(t) - c_3 \Gamma(t) - c_4 \gamma_0 \Gamma(t), \\
 w(t) &= c_1 \frac{N+1}{N} - c_2 \Gamma^2(t) + c_4 \gamma_0 (N+1) \Gamma(t), \\
 u(t) &= c_5 \Gamma(t), \\
 v(t) &= c_6 \Gamma(t)
 \end{aligned} \tag{11}$$

(ii) $N = 0$

$$\begin{aligned}
 x(t) &= d_4 \Upsilon^2(t), \\
 y(t) &= d_2 \Upsilon(t) + d_3 \Upsilon(t) - d_4 \Upsilon^2(t), \\
 z(t) &= -d_3 \Upsilon(t) - d_4 \Upsilon^2(t), \\
 w(t) &= d_1 - d_2 \Upsilon(t) + d_4 \Upsilon^2(t), \\
 u(t) &= d_5 \Upsilon(t), \\
 v(t) &= d_6 \Upsilon(t)
 \end{aligned} \tag{12}$$

where $\Gamma(t) = e^{-(1+2N)\gamma_0 t}$, $\Upsilon(t) = e^{-\gamma_0 t}$, and the coefficients $c_1, c_2, c_3, c_4, c_5, c_6$ in Eq. (11), and $d_1, d_2, d_3, d_4, d_5, d_6$ in Eq. (12) are determined by the corresponding initial conditions.

4. Entanglement dynamics with the initial conditions

4.1. $N > 0$

We now analyze the entanglement dynamics. Starting from the initial states $\rho_{AB}^\Phi(0)$ in Eq. (7), the coefficients of Eq.(11) are determined as

$$\begin{aligned}
 c_1 &= \frac{N(N+1)}{(2N+1)^2}, \\
 c_2 &= \frac{N(N+1)}{(2N+1)^2} - \frac{1-r}{4}, \\
 c_3 &= \frac{1}{2(2N+1)^2} + \frac{r}{2}(\cos^2(\alpha) - \sin^2(\alpha)), \\
 c_4 &= -\frac{1}{\gamma_0(2N+1)^2}, \\
 c_5 &= r \sin(\alpha) \cos(\alpha), \\
 c_6 &= 0
 \end{aligned} \tag{13}$$

Then the concurrence of $\rho_{AB}^\Phi(t)$ is

$$C(\rho_{AB}^\Phi(t)) = 2 \max\{0, |u(t)| - \sqrt{x(t)w(t)}\}. \tag{14}$$

So the ESD appears when,

$$|u(t)| - \sqrt{x(t)w(t)} \leq 0 \Leftrightarrow u^2(t) - x(t)w(t) \leq 0. \tag{15}$$

Obviously,

$$t \rightarrow +\infty, \quad u^2(t) - x(t)w(t) = -\frac{N^2(N+1)^2}{(2N+1)^4} < 0,$$

and

$$t = 0, \quad u^2(t) - x(t)w(t) = r^2 \sin^2(\alpha) \cos^2(\alpha) - \frac{(1-r)^2}{16}. \tag{16}$$




Figure 2. (Color online) ($r = 1, \alpha = \pi/4$) Plot of concurrence of $\mathcal{C}_\rho^\Phi(t)$ vs “N” and $\gamma_0 t$.

Thus, ESD always occurs if the initial state is entangled, which means $\frac{1}{1+2|\sin 2\alpha|} \leq r \leq 1$, according to Peres’ criterion.

Note that Ref. [23] found that when the mean thermal photon number is not zero, in the thermal reservoir the entanglement sudden death always happen from the simulation. However here we get the same result from the theoretic analysis. In Fig. 2, 3, and 4 we simulate the concurrence as a function of $\gamma_0 t$ and one of the parameters N, α , and r , respectively. Fig. 2 is the concurrence $\mathcal{C}_\rho^\Phi(t)$ as function of $\gamma_0 t$ and N , fixing the purity $r = 1$ and initial degree of entanglement $\alpha = \frac{\pi}{4}$, the Bell-like states. It shows that ESD time is affected by N . The smaller N , the longer its ESD time. In Fig. 3, we plot the concurrence $\mathcal{C}_\rho^\Phi(t)$ as function of $\gamma_0 t$ and initial entanglement α when $r = 1$. When $\alpha = \frac{\pi}{4}$ or $\alpha = \frac{3\pi}{4}$ the initial states are reduced to Bell-like states. The ESD time is sensibly affected by α . Fig. 4 is the concurrence as function of $\gamma_0 t$ and purity r . As exemplified before, when $0 \leq r \leq \frac{1}{1+2|\sin(2\alpha)|}$ the initial state is separated, so we choose r from $1/3$ to 1 . It shows that the larger the purity r the longer the ESD time.

Here we study the entanglement dynamics of bipartite quantum system in the global environment effect. However, Ref. [26] had shown that under the classical noise effect entanglement may experience a sudden death process even if the local coherence of one participating particle is well preserved and the other one decays to zero asymptotically. How do the local thermal reservoirs influence the entanglement dynamics? Up to my knowledge, the master equation needs to be reconstructed and the local environment effect embodied by the spectral density of the thermal reservoir $J(\omega, T)$. The one-body decoherence dynamics was studied in [12, 33] under the local environment effect. Whether does the multipartite entanglement dynamics under the local thermal reservoirs hold under the classical noise effect proved by Yu et al [26]? We will study it in our further work.




Figure 3. (Color online)($r = 1, N = 0.1$)Plot of concurrence of $\mathcal{C}_\rho^\Phi(t)$ vs α and $\gamma_0 t$.




Figure 4. (Color online)($\alpha = \pi/4, N = 0.25$)Plot of concurrence $\mathcal{C}_\rho^\Phi(t)$ vs “r” and $\gamma_0 t$.

4.2. $N = 0$

The coefficients in Eq. (12) have the form

$$\begin{aligned}
 d_1 &= 1, \\
 d_2 &= 1, \\
 d_3 &= -\frac{1-r}{2} - r \sin^2(\alpha), \\
 d_4 &= \frac{1-r}{4}, \\
 d_5 &= r \sin(\alpha) \cos(\alpha), \\
 d_6 &= 0.
 \end{aligned} \tag{17}$$

Thus

$$\begin{aligned}
 &u^2(t) - x(t)w(t) \leq 0 \\
 \Leftrightarrow &\Upsilon^2(t) - \frac{4}{1-r}\Upsilon(t) + \frac{4}{1-r} - \frac{16r^2}{(1-r)^2} \sin^2(\alpha) \cos(\alpha) \geq 0.
 \end{aligned} \tag{18}$$

So the ESD occurs when,

$$\frac{1}{1 + 2|\sin(2\alpha)|} < r < \frac{-1 + \sqrt{1 + 4\sin^2(2\alpha)}}{2\sin^2(2\alpha)}, \quad (19)$$

and the ESD time is

$$\gamma_0 t^* = \ln(1 - r) - \ln[2(1 - \sqrt{r + r^2 \sin^2(2\alpha)})]. \quad (20)$$

When $\gamma_0 t \in [\gamma_0 t^*, +\infty)$, the concurrence $C_{AB}^\Phi = 0$. In Fig. 5 we plot the ESD area. If the initial state is pure, i.e., $r = 1$, we find that $x(t) \equiv 0$. Then the concurrence is $C(\rho_{AB}^\Phi(t)) = 2|u(t)| = |\sin(2\alpha)|e^{-\gamma_0 t}$, which implies that the nonexistence of ESD. After we submitted our paper, a closely related work appeared in [28]. Ikram et al extended their former result [23] to more general state for the zero temperature limit and provided some discussions on the thermal case. Comparing with us, there are three main differences. Firstly, although both papers studied the same system, a two 2-level atom system, different initial states had been chosen. These two initial states are independent, and the results are complementary. Secondly, for pure 2-qubit entangled states in a thermal environment, Ref. [28] says that “We see in plots of figure 3 that entanglement sudden death always happens for non-zero average photon number in the two cavities and entanglement sudden death time depends upon the initial preparation of the entangled states.” Here, a theoretical analysis is provided by Eqs. (14, 15, 16). Finally, in the vacuum reservoir we give the sufficient and necessary condition Eq.(19) for the entanglement sudden death, and answer the question when the ESD occurs and what ESD time is. Furthermore, the ESD area relies on the initial purity r and initial entanglement α , see Fig. 5. Here we give a more detail examination and analysis.

The ESD condition and corresponding time for initial state $\rho_{AB}^\Phi(0)$ at $N = 0$ are summarized in the following table.

TABLE I. ESD condition and time for initial state $\rho_{AB}^\Phi(0)$ at $N = 0$

α	$(0, 2\pi)$
Condition	$\frac{1}{1+2 \sin(2\alpha) } < r < \frac{-1+\sqrt{1+4\sin^2(2\alpha)}}{2\sin^2(2\alpha)}$
Time	$\gamma_0 t^* = \ln(1 - r) - \ln[2(1 - \sqrt{r + r^2 \sin^2(2\alpha)})]$

5. Conclusions

In summary, we have analyzed the interesting phenomenon of entanglement sudden death determined by the dimensionless parameters $\hbar\omega_0/k_B T$, α , and r . Sufficient conditions for ESD have been given for initial state $\rho_{AB}^\Phi(0)$. We examine the conditions for ESD and calculate the corresponding ESD time by the Wootters’ concurrence. We observe that ESD is determined by the parameters like N , amount of initial entanglement α , and the purity r . For $N = 0$, we get the analytical expressions of the ESD condition and ESD time. For $N > 0$ we give a theoretical analysis that ESD always occurs, and simulates the concurrence as a function of $\gamma_0 t$ and one of the parameters




Figure 5. (Color online) Plot the ESD area of the extended Werner-like states $\rho_{AB}^{\Phi}(t)$ with $N = 0$ for $\alpha \in [0, 2\pi]$. The red line is $r = \frac{-1 + \sqrt{1 + 4 \sin^2(2\alpha)}}{2 \sin^2(2\alpha)}$, and the blue line is $r = \frac{1}{1 + 2|\sin(2\alpha)|}$.

N , α , and r . In fact, these above results can also be obtain for other “X” initial states, for example $\rho_{AB}^{\Psi}(0) = r|\Psi\rangle_{ABAB}\langle\Psi| + \frac{1-r}{4}\mathbf{I}_{AB}$, where $\Psi_{AB} = (\cos(\alpha)|00\rangle + \sin(\alpha)|11\rangle)_{AB}$.

We analyze the dynamic behavior of entanglement in the open quantum system But our ultimate aim is to control the entanglement such that it be a resource for practical realization. Despite of the noticeable progresses of entanglement, many fundamental difficulties still remain. One of the problem is ESD due to the interactions between system and environment; the other is that as the N -particle increased, the entanglement becomes arbitrarily small, and therefore useless as a resource [34]. Here we indicate that preparing some initial states can help prolong the ESD time. However, we think that it is a long way how to design some effective control field to make the N -particle large enough and protect the entanglement that make it practically useful.

Acknowledgments

We thank the referees for improving the manuscript. This research is supported by the National Natural Science Foundation of China (No. 60774099, No. 60221301) and by the Chinese Academy of Sciences (KJCX3-SYW-S01).

References

- [1] Bell J S 1964 *Physics* **1** 195
- [2] Einstein A, Podolsky B and Rosen R 1935 *Phys. Rev.* **47** 777
- [3] Nielsen M A and Chuang I L 2000 *Quantum Computation and Quantum Information* (Cambridge: Cambridge University Press)
- [4] Bennett C H and DiVincenzo D P 2000 *Nature(London)* **404** 247
- [5] Bennett C H, Brassard G, Crepeau C, Jozsa R, Peres A and Wootters W K 1993 *Phys. Rev. Lett.* **70** 1895
- [6] Bouwmeester D, Pan J W, Weinfurter M, and Zeilinger A 1997 *Nature(London)* **390** 575

- [7] Ekert A K 1991 *Phys. Rev. Lett.* **67** 661
- [8] Breuer H P and Petruccione F 2002 *The Theory of Open Quantum Systems* (Oxford: Oxford University Press)
- [9] Weiss U 1999 *Quantum Dissipative Systems (Second Edition)* (Singapore: World Scientific Publishing)
- [10] Mintert F, *et al.* 2005 *Physics Reports* **415** 207
- [11] Branderhorst M P A, *et al.* 2008 *Science* **320**, 638
- [12] Cui W, Xi Z R and Pan Y 2008 *Phys. Rev. A* **77** 032117
- [13] Zhang J, Li C W, Wu R B, Tarn T J, and Liu X S 2005 *J. Phys. A: Math. Gen.* **38** 6587
- [14] Yu T and Eberly J H 2004 *Phys. Rev. Lett.* **93** 140404
- [15] Yu T and Eberly J H 2006 *Phys. Rev. Lett.* **97** 140403
- [16] Almeida M P, *et al.* 2007 *Science* **316** 579
- [17] Laurat J, *et al.* 2007 *Phys. Rev. Lett.* **99** 180504
- [18] Wang J, *et al.* 2006 *J. Phys. B* **39** 4343
- [19] Dajka J, Mierzejewski M and Łuczka J 2008 *Phys. Rev. A* **77** 042316
- [20] Dajka J, and Łuczka J 2008 *Phys. Rev. A* **77** 062303
- [21] Roos C 2008 *Nature Physics* **4** 97
- [22] Konrad T, *et al.* 2008 *Nature Physics* **4** 99
- [23] Ikram M, Li F L , and Zubairy M S 2007 *Phys. Rev. A* **75** 062336
- [24] Al-Qasimi A and James Daniel F V 2008 *Phys. Rev. A* **77**, 012117
- [25] Huang J H and Zhu S Y 2008 *Opt. Comm.* **281**, 2156
- [26] Yu T and Eberly J H 2006 *Opt. Comm.* **264** 393
- [27] Huang J H and Zhu S Y 2007 *Phys. Rev. A* **76** 062322
- [28] Tahir R, Ikram M, Aazim T, and Zubairy M 2008 *J. Phys. B* **41** 062332
- [29] Wootters W K 1998 *Phys. Rev. Lett.* **80** 2245
- [30] Turchette Q A, Myatt C J, King B E, Sackett C A, Kielpinski D, Itano W M, Monroe C and Wineland D J 2000 *Phys. Rev. A* **62** 053807
- [31] Maniscalco S and Petruccione F 2006 *Phys. Rev. A* **73**, 012111
- [32] Peres A 1996 *Phys. Rev. Lett.* **77** 1413
- [33] Villares Ferrer A, and Morais Smith C 2007 *Phys. Rev. B* **76** 214304
- [34] Aolita L, Chaves R, Cavalcanti, Acín A, and Davidovich L 2008 *Phys. Rev. Lett.* **100** 080501